

The **763C>G** Polymorphism of The Secretory **PLA2IIa** Gene is Associated with Endometriosis in Iranian Women

Mehdi Sahmani, Ph.D.¹, Masoud Darabi, Ph.D.², Maryam Darabi, Ph.D.¹, Talaat Dabaghi, M.D.³,
Safar Ali Alizadeh, M.Sc.¹, Reza Najafipour, Ph.D.^{1*}

1. Department of Clinical Biochemistry, Cellular and Molecular Research Center, Qazvin University of Medical Sciences, Qazvin, Iran
2. Department of Biochemistry and Clinical Laboratories, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran
3. Kosar Hospital, Qazvin, Iran

Abstract

Background: Endometriosis is a chronic gynecological disease resulting from complex interactions between genetic, hormonal, environmental and oxidative stress and intrinsic inflammatory components. The aim of this study was to investigate the potential association of the 763C>G polymorphism in the secretory phospholipase A2 group IIa gene (PLA2G2A) with the risk of endometriosis in Iranian women.

Materials and Methods: Ninety seven patients with endometriosis along with 107 women who were negative for endometriosis after laparoscopy and laparotomy, and served as the control group, were enrolled for this cross-sectional study. Samples were genotyped using the polymerase chain reaction-restriction fragment length polymorphism method.

Results: Multivariate analysis was used to examine the association between the risk of endometriosis and the 763C>G polymorphism of PLA2G2A. Genotype distributions of PLA2G2A were significantly different between patients and the controls ($p<0.001$, OR=0.22, 95% CI=0.21-0.39). Correlation analysis showed that there was a significant association between the normal homozygous genotype and susceptibility to endometriosis ($p<0.001$).

Conclusion: The present study suggests that the 763C>G polymorphism of PLA2G2A plays an important role as an independent factor in the risk of endometriosis in Iranian women.

Keywords: Endometriosis, polymorphism genetic, PLA2G2A

Citation: Sahmani M, Darabi M, Darabi M, Dabaghi T, Alizadeh SA, Najafipour R. The 763C>G polymorphism of the secretory PLA2IIa gene is associated with endometriosis in Iranian women. *Int J Fertil Steril*. 2015; 8(4): 437-444.

Introduction

Endometriosis is one of the most frequent disorders in gynecology that is characterized by development of endometrial tissue outside the uterus (1). Endometriosis affects >10% of all women of reproductive age and >30% of all infertile women (2, 3). The common symptoms of endometriosis are pelvic pain, dysmenorrhoea, dyspareunia and infertility (4). The disease is diagnosed by laparoscopy with or without biopsy for histological diag-

nosis (4, 5). According to extent, endometriosis is classified as stage I (minimal), stage II (mild), stage III (moderate) and stage IV (severe) (6). Previous studies have demonstrated that oxidative stress and inflammatory activity as well as genetic abnormalities may be associated with the development and progression of endometriosis (2, 7, 8).

Similar to the process that exists in cardiovascular disease, an abnormal lipid profile including increased

Received: 03 Jul 2013, Accepted: 28 Jan 2014

* Corresponding Address: Department of Clinical Biochemistry, Cellular and Molecular Research Center, Qazvin University of Medical Sciences, Qazvin, Iran

Email: rnajafipour@gmail.com



Royan Institute
International Journal of Fertility and Sterility
Vol 8, No 4, Jan-Mar 2015, Pages: 437-444